4/	1
11/01	1/nn
ver	100/7/8
4	タ/フ/れ
	9/1/0

1							NA	2/1/8/71	8/7/80	
Shorebard ,	record	o- (	oral	velle	Reservoin	. summe	(1980)	9/1/	4	
	654	12 July	ام تدار	26 Tuly	308					
Simi. Plover	-	2	-	8	6		P			
Kildeer	17	18	25	8	6					
Solit. sandp.	3	3		5	1					
Spot. Sandp.	3	10	3	14	8					
l Eyellowkys	2	26	26	52	52					
gr. jellowlegs	_	_	-	8,	2					
poect. sandp.	_	+	13	57	39					
Ucst Sondp.		70	37	66	38					
Semipal soudp.	-	1	2	28	15			,		
still sadge.		_	, q	-	-					
1-6 downthen	_	-	-	_	ı					
S-b. dowitcher	_	8,	8	_						
surdp., sp.	_	8	2	_	1					
1/4										
	i.all a	el.			4. 1mm.b.	id same budy	~?			
	2. of bull- and: 2i, uad				5: same birds					

2. of bull og i: 2i, 11acl

3. all in remains of br. pl.

locations: Sand Pout, ponds to e. + w. of O-Roca, ponds s. of Con Falls Rd, to s. of farm between 218+0-RD.

Notes on dowitchers, summer 1980

short-billed (8); 12-19 July, Coralville Res., pond west of O-road.

ochre breast, white belly; spotting on side of breast in vicinity of bend of folded wing; very conspicuous pale feather edging on back, scapulars, KNYEKER tertials. Associating with lesser yellowlegs, stilt sandpipers. Resting, preening, feeding in drying pool. Fed belly-deep in water.

short-billed, imm. 1 27 July Cone Marsh.

breast bright ochre-buff, unmarked except for a grey smudge at each side of the base of neck; belly and undertail coverts white. Face buffy grey, dark mark through eye, distinct paler supercilium. crown brwonish, not so dark as back. Back and wing-coverts dark, with prominent pale margins to feathers. tertials prominently banded with dark and pale bars or blotches, hars dark and light bars/blotches of equal width, uniformly spaced along feathers. bill dark, legs dull greenish-orange. resting/feeding preening on shore of small mud-rimmed pool. fed on mud, not in water. no other shorebirds present.

long-billed, 1 ad 3 Aug. Coralville Res., pond on north side.

breast and underparts (to undertail coverts) deep rufous-buff; undertail coverts buffy, with short bars near the tips of each feather. Strong barring at side of breast. Some barring on flanks. Scapulars dark with narrow pale edges, teritals dark brownish/black with crossed with narrow pale bars. Wing-coverts mixture of medium-grey feathers and feathers with dark centers and narrow pale edges. No spotting on central portion of breast or on belly. Tail with black and light bars, of approximately equal width. Bill dark, very long, obvious at first glance as longer than that of the short-bills seen previously. Legs pale yellowish-green Face grey, with a dqrker mark thru eye and a paler superciliary stripe. Crown dark brown, hind neck a slightly paler, greyer brown. On a pond with a narrow border of mud, then tall weeds. Did not associate with the lesser yellowlegs, pectoral sandpipers, EXXREED least or semipalmated sandpipers present. Fed with head down, steady motion of bill, in water to tarsal joint.

Other records, summer 1980

redtailed hawk: 1 imm Swan Lake area 19 July and 3 Aug; 1 imm Cone marsh 27 July.

turkey vulture 3 Coralville Res 19 July

black tern 1 Coralville Res 19 JUly

Sterna sp, prob. Forster's: 8 Coralville Res 26 July.

hairy woodpecker l Iowa City 4 August.