

United States Department of the Interior

FISH AND WILDLIFE SERVICE DESOTO NATIONAL WILDLIFE REFUGE RR 1, BOX 114 MISSOURI VALLEY, IOWA 51555 (712) 642-4121

September 23, 1981

Dr. Jim Dinsmore
Department of Animal Ecology
Science Hall II
Iowa State University
Ames, IA 50011

Dear Dr. Dinsmore:

On August 8, 1981, two adult and one immature least tern (<u>\$terna albifrons</u>) were sighted at California Bend Wildlife Managment Area (ICC). They were present on a sandy area exposed by low water conditions at the east edge of the outlet (marked by X on attached map). The other observer and his current address is:

Ken Grannemen Squaw Creek NWR P.O. Box 101 Mound City, Missouri 64470

Least terms have also been intermittently sighted at DeSoto NWR throughout summer 1981. We have not kept record whether they were mature or immatures, however. Although my memory tells me I saw immatures, I cannot verify it. We do not know of any least term nesting attempts on the refuge in 1981.

Thanks for the information you sent.

Sincerely,

Rick

Rick Sojda Assistant Refuge Manager

Enclosure

	DOCUMENTATION FORM for extraordinary bird signtings in lowa
	What species? Prairie Folcon How many?
	Location? De Soto National Wildlife Refuge
	Type of habitat? Open Country, Cropfields, Scattered trees
	When? date(s): Nov. 8, 1981 time: 9 AM to
	Who?your name and address: ERWIN E. KLAAS
	others with you: Ben KLAAS
	others before or after you:
	Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, under tail, legs, feet). Also mention voice and behavior.
	Obviously, a large falcon (Crow Size) with pointed wings and long narrow tail. Bird was seen in flight moving slowly
	into strong headwind at about 30 yards. Upper parts were
	brown or chestnut and facial morkings were clearly seen.
	Process of dork axillar markings distinguished it from
	Presence of dark axillor markings distinguished it from immature peregrine falcon. Underports were light with a few
0	Hark Streaks.
	Similar species and how eliminated: See above.
	Did any one disagree or have reservations about identification?
	If yes, explain: has been sighted Several times this Fall on refuge.
	Pright Sunlight with Sun behind observer. Distance estimated
	Previous experience with species and similar ones: Have Seen both in the wild, including Peregrape Falson only 2 weeks
	References and persons consulted before writing description. None
	How long before field notes made?this form completed?thrs
	MAIL TO: T. H. Kent, Field Reports Editor, 211 Richards Street, Iowa City IA 52240