Records Committee, Iowa Ornithologists' Union

Barrow's Goldeneye 28 Nov 1994 Tera L., Johnston, Polk Co., IA Reid and Pam Allen P-0434, IBL 65:12, 81 Record Number: 94-46 Classification: A-P

DOCUMENTATION Reid and Pam Allen, 1601 Pleasant St., West Des Moines 50265 [6/26/95*] PHOTOGRAPH Reid Allen, P-0434 REFERENCES Field Reports: IBL 65:12 Records Committee: IBL 65:81 VOTE: 7 A-P A-P, Excellent photo vs Common Goldeneye; "forehead" shape clearly diagnostic.



Barrow's Goldeneye Johnston, Polk Co., IA 28 Nov 1994 Reid Allen 94-46 P-0434

DOCUMENTATION

94-46

Barrow's Goldeneye (Bucephala islandica) SPECIES: HOW MANY: 1 Tera Lake, IPTV Pond, Johnston, Polk County, Iowa OCATION: Enclosed freshwater pond. HABITAT: 28 November, 1994 DATE: TIME & LENGTH OF 12:15 - 12:20 p.m. OBSERVATION: FIRST OBSERVATION Jerry Jancik on the day before MADE BY: Reid & Pam Allen, 1601 Pleasant Street, DOCUMENTER'S NAME & West Des Moines, IA 50265 ADDRESS: Eloise and Eugene Armstrong; Maridel Jackson OTHERS WHO SAW BIRD: The head shape was the most noticeable characteristic which differentiated the Barrow's DESCRIPTION OF BIRD: Goldeneye from the Common Goldeneyes. The forehead of the Barrow's Goldeneye was very steep and almost straight up and down, compared to the Common Goldeneve whose forehead was gently sloped from the top of the bill to the rounded head. The Barrow's Goldeneve crescent white patch on either side of the cheek, below the eye, was variable on either side of the face, although it was very definitely a crescent-shaped white patch, as opposed to the rounded white patch on the Common Goldeneyes. The bill on the Barrow's was short and stubby, appearing black. The eye was a bright golden yellow. The head was black, with a slight very dark purplish cast to it. The breast and sides were white, with a white collar extending around the neck. The sides of the wings looked like a series of white "port holes" or "windows" on each side of the bird and it looked as if there was a black hook extending down the side of the bird at the front end of the wing into the breast area. The back and tail were black, and the black on the sides of the body seemed to extend past the flank area toward the front of the bird. The tail was short and the whole body of the duck sloped down to the water. The Barrow's seemed about the same size as the Common Goldeneyes. Common Goldeneve eliminated because of head shape, wing and side markings, facial SIMILAR SPECIES AND pattern. HOW ELIMINATED: Clear day, but very windy. VIEWING CONDITIONS: PREVIOUS EXPERIENCE Viewed at DeSoto Bend NWR in Iowa a few years ago. WITH SPECIES: Kowa TSN-4 spotting scope; Bushnell 8x36 Custom Binoculars. EQUIPMENT: Minolta X-700 camera. FIELD GUIDE TO THE BIRDS OF NORTH AMERICA, National Geographic Society, Second REFERENCES CONSULTED: Edition, 1987. HOW LONG BEFORE Immediately upon leaving sighting. FIELD NOTES MADE: HOW LONG BEFORE THIS January 31, 1994. FORM COMPLETED: Photograph (1) enclosed taken by Reid I. Allen on November 28, 1994. PHOTOGRAPH: