## IOWA BREEDING BIRD ATLAS DOCUMENTATION FORM

What species?Northern Goshawk	How many?
Location:County Boone	How many?1 
Location in detail Boone Wildlife	Research Station.just south of Ledges
State Park	adult near the Des Mulnes River and flushed one ad
Habitat?Oak woods, open fiel	ds
	Time: 1:00 pm to to second and a second
	shat the species press of the block of the constant of the con
	Road, Boone IA 50036

Description of bird(s): include size, shape, detailed description of all parts including bill, eye, head, neck, back, wings, tail, throat, breast, belly, undertail, legs, feet; also voice and behavior. Use additional sheets if necessary.

Bird was observed flying overhead at treetop height. It flew with a rather characteristic flap-glide pattern of accipiter hawks, which is what caught my eye in the first place. It did soar for a short distance after flying overhead. The bird was large for a hawk and was gray underneath (suggesting an adult) with a banded tail and a lighter neck-head area. The wings were relatively large (chunky) and one primary feather was missing from the right wing.

If books or other birders were consulted before writing up the documentation, please list them here:

Other observers were DeWarne Jackson, Wijdlite Biologist, iom rozham, Fisr Biologist, and Jim Kienzler, Wildlife Biologist. Ne Juoble-chucked with J Peterson Field Bidde to Birds of Eastern North America.

Similar species and how eliminated: Too large for a Cooper's hawk (it was big!!), or a broad-winged hawk. No other gray hawks of that size and description that I am aware of.

Did anyone have reservations about identification? If so, explain: Because of the time of year, we had reservations but we rather methodically eliminated other possibilities based on behavior and size. Viewing conditions: lighting, distance and how measured, optical equipment used.

Clear sky, bird observed with unaided eye at approximately 30 yards straight overhead (treetop height). Sun was at our backs, bird in front.

## Previous experience with this species and other similar ones:

We have an immature goshawk at the station presently. I have photographed an adult near the Des Moines River and flushed one adult off a bobwhite quail it had killed (McCoy Wildlife Area)

BREEDING BEHAVIOR: Give all details observed that indicate that the species breeds on the block. If nest, eggs, young, or courtship display were observed, describe in detail.

Not Applicable.

Description of bird(s): include size, shape, detailed description of all parts including bil), hyp. head, neck, back, winge, tail, throat, broast, bally, undertail, lage, fast! also voice and berevior. Use additional shapes if necessary.

Utro was observed Flying overhead at treetop height. It flew with a rather characteristic Flap-olide pattern of accipiter hawks, which is what caught by eye in the first place. It did soar for a short distance after flying overhood. The bird was large for a hawk and was gray underneath (succesting an adult) with a banded tail and a lighter neck-head area. The wings were relatively large (chunky) and one primary feather was missing from the right wing.

If books or other birders were consulted before writing up the documentation, please list them here:

Other observers were DeWaine Jackson, Wildlife Biologist, Tom Putnam, Fisheries Biologist, and Jim Kienzler, Wildlife Biologist. We double-checked with a Peterson Field Guide to Birds of Eastern North America.

How long after observation were field notes made?

Immediately and the state of the same and the same and on the same beauty beauty beauty

How long after observation was this form completed? 2 hours

> Jecruse of the time of year, we had reservations but we rather methodically illutrated other possibilities based on behavior and size.