

UNIVERSITY OF NORTHERN IOWA

03-18-92 Jim Thank you for the documentation form. The made copies and hope to use them in the Future. From the information I've seen there is no earlier date for a Barn Swallow sighting. Can you let me know if this is correct. I'd like to know if I set a new record. I spoke to the biologists at other Creek and 3 Sandhill Cranes spent last summer there. I am going to be working for the DNR at Otter Creek this summer so I'll be up on all the activity there and will let you Know. So far no sight of them yet this year. I'll be in touch Thanks Again CSA Inc. DSN. 1038 Printed in USA College Letters over->

UNIVERSITY OF NORTHERN IOWA

PS. IF your ever up this way let me know, I'd like to meet you and possibly go birding together. Nik

\$ C5A Inc. 08N, 1924, Pr. ---- --- -----

DOCUMENTATION FORM for extraordinary bird sightings in Iowa		
What species?	Baco Swallow	How many? One
Location? Section #7, Otter Creek Marsh, Tama County, Iawa .		
Type of habitat? Cattail Marsh		
When? date(s): March 5, 1992 time: 14:00 to 14:30		
Who?your name and address: Mike Stegmann 1165 260th Street, Garwin Fowa others with you:		
others with yo	u:	50632
others before	or after you:	

Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, under tail, legs, feet). Also mention voice and behavior.

While observing waterfowl in the unusually warm weather (temperature in the upper 60's) I noticed a Swallow type bird which Flew through my field of vision. The flight was the first cluelethe birds identification. Flight was constant Swooping and close to waters surface much like TreeSwallows. Color of the bird was blue-black upper side (wings, tail, back, head) and light tan underside (breast, belly). Also noted was a chin strap and dark amber or honey color throat. The shape was that of a swallow and the tail was long and deeply forked. The bird was abserved for approximately 30 minutes and flight was continues in the same general area, about a 200 yard circle.

Similar species and how eliminated: Tree Swallows USD this area to a great extent later in the spring and summer but the bird seen was blue-black with tan underside and long forked tail which tree swallows lack. Did any one disagree or have reservations about identification? No

If yes, explain:

Viewing conditions: give lighting, distance (how measured), and optical equipment: Lighting was good considering overcast skits with sprinkles. Distance was estemated at 50 to 100 yards. Tasco brand binoculars with zoom 7x-35x50 power were used.

Previous experience with species and similar ones: Very familiar with species. Nests on my property.

References and persons consulted before writing description: Jim Fuller

How long before field notes made? Immediately this form completed? 12 Days

MAIL TO: T. H. Kent, Field Reports Editor, 211 Richards Street, Iowa City IA 52240