

Ferruginous Hawk

< 1907

Sioux City, Woodbury Co., IA

Rich

Anderson 1907, Bailey 1918

Record Number: 93-GK

Classification: NA

REFERENCES

Anderson 1907

Bailey 1918

VOTE: 5 NA

NA, Rich saw some mounted birds in stores but did not verify  
their place of origin./thk

## Ferruginous Hawk

151. (347a). *Archibutea ferrugineus* (Licht.). Ferruginous Rough-leg.

Anderson 1907  
p 252

The Ferruginous Rough-leg occurs occasionally in Iowa during the migrations. It is a bird of the western United States, occasionally straggling east to Iowa or even to Illinois and Wisconsin. Major Bendire states (Life Hist. of N. A. Birds, 259-60) that it "has been reported as nesting near Grinnell, but the record has not been fully verified." Mr. Lynds Jones says of this: "I know nothing about the Ferruginous Rough-leg at Grinnell. Bendire concluded that the Grinnell bird must be Ferruginous, and so stated upon his own motion. I always questioned it." In Nebraska it is "the common Rough-legged Hawk in the state, and occurs throughout. It is less common in the eastern portion" (Rev. Bds. Neb., 1904, 52).

County records: Blackhawk—"rare winter visitant; specimen in museum of I. S. N. S." (Walters). Kossuth—"a few seen. I am sure of identity of this species, as it is well known to me" (Bingaman). Linn—"a pair of Ferruginous Rough-legs winter every year at Gordon's grove, Waubec; have seen them there four different years. They are larger birds than the Swainson, but are white or nearly so below, with feathered legs. Mr. J. W. Preston collected a set of Ferruginous Rough-leg near Newton" (Berry). Pottawattamie—"straggler" (Trostler). Woodbury—"rare; it is not uncommon in the various mounted Hawks seen in the stores of Sioux City. I have never handled a fresh specimen" (Rich).

Bailey, B. H. 1918. The Raptorial Birds of Iowa, Iowa Geological Survey Bull. Des Moines pp 139-141




FIG. 55.—Ferruginous Rough-legged Hawk.

Subgenus Brewsteria Maynard.

15. *ARCHIBUTEO FERRUGINEUS* (Lichtenstein). 348.  
Ferruginous Rough-leg. California Squirrel Hawk.

*Field Characters.*—A large reddish backed, white breasted hawk. Sluggish as compared with other hawks, and found hunting early and late in open fields.

*Description.*—Above, rufous, especially on the wings, the centers of the feathers dusky or brownish black; tail ashy white, the feathers edged with tawny. Below, white, with a few reddish streaks on the sides of upper breast and body. Legs sharply contrasting with other parts below, being reddish with numerous black bars, and feathered to the tocs.

No.	SEX	LOCALITY	DATE	BIRDS	MAMMALS	MISCELLANEOUS	NOTES
1	Male	Iowa					Empty
2	Female	Osage	Dec. 27, 14				Empty
3	Male	Britt	Dec. 24, 14		Small mammal		
4	Male	Iowa	Nov. 23, 14		Small mammal		
5	Male	Marion Co.	Nov. 6, '05		2 shrews—3 rodents	Several grasshoppers	
6	Male	(Glendale)	Nov. 10, '05				Empty
7	Male	Boone	Sep. 1, '06		2 frogs.		
8	Male	Iowa	—, '06				Empty
9	Male	Cass	Nov. 10, '06				Empty
10	Male	Strait Center	Jan. 4, '06		2 mice		
11	Male	Iowa	May 1, '07		Hair		
12	Male	Gladbrook	Nov. 25, '07		Several mice.		
13	Male	Reinbeck	Nov. 13, '07				

LIST OF STOMACH CONTENTS




FIG. 66.—Map showing the distribution in Iowa of the Perrenigous Hawk.

- Hedley, H., Two hundred Wild Birds of Iowa, p. 62, 1906.  
Widmann, Otto, A Preliminary Catalogue of the Birds of Missouri; Trans. Acad. Sci., St. Louis, pp. 99-100, 1907.  
Anderson, R. M., Birds of Iowa, p. 262, 1907.  
Gory, C. H., Birds of Illinois and Wisconsin, p. 469, 1909.  
A. O. U. Check List, p. 161, 1910.  
Chapman, F. M., Birds of Eastern North America, p. 300, 1912.

- Keeges, Charles, Hawks and Owls of the U. S., p. 91, 1893.  
Fisher, A. K., Hawks and Owls of the U. S., p. 17, April, 1897.  
Hedley, Capt. Chas., Life Histories of North American Birds, Vol. I, Pt. 259, 1892.  
Hatch, Dr. J. L., Birds of Minnesota, p. 194, 1892.  
\*Coates, E., Key to North American Birds, 4th Edition, p. 551, 1892.  
Hedge, Vol. 16, p. 132, 1891.  
\*Kelsay, Carl, Birds of Poweshiek County, Iowa: Ornithologist and Otolologist, N. S., History of the Birds of Kansas, p. 271, 1881.  
Hedgeway, R., Ornithology of Illinois, p. 481, 1889.  
Jones, L., and Parker, H. W., in W. W. Cooke's Report on the Bird Population in the Mississippi Valley, 1884-85; U. S. Department of Agriculture, Division of Economic Ornithology, Bull. 2, p. 117.  
\*Jones, L., and Parker, H. W., in W. W. Cooke's Report on the Bird Population in the Mississippi Valley, 1884-85; U. S. Department of Agriculture, Division of Economic Ornithology, Bull. 2, p. 117.  
\*Coates, E., Key to North American Birds, 3d Edition, p. 551, 1887.  
Archibald, Ferruginous (Lichenalelm).

Although, because of their infrequent visits to the state they are mainly food is mostly small mammals such as ground squirrels, field mice, and rabbits, and also reptiles.  
Their flight simulates that of the Osprey.  
By those who have watched these birds closely, their flight when soaring at a great height is described as resembling that of the Golden Eagle; and when winging the air a short distance above the ground their flight simulates that of the Osprey.

1907.)  
Several authors have referred to it as nesting near Grinnell, but erroneously attributed the record. (Anderson, Birds of Iowa, p. 252,  
this record is denied by Mr. Lynds Jones, to whom Captain Benard  
corroborated the record.

Jalls.  
Iowa is in the collection of the State Teachers' College at Cedar Rapids in as the "handsomest of the North American Falcoinidae."  
The only specimen which I have ever seen that has been taken in the east of the Mississippi, Dr. Elliott Coates is said to have re-taken and comes only occasionally into Iowa, though it is sometimes seen in the open prairies of the west and composes only a small part of the bird life of the state.  
This large hawk is conspicuously a bird of the open prairies of the cornish and Illinois.

Many.—Western United States, wandering casually to Iowa, Wisconsin and Illinois.  
tail, 10 to 11 inches; length, 22 to 24 inches; wings, 16 to 18 inches;  
11 to 12 inches.