

Chestnut-collared Longspur

CA 1892

Cedar Rapids, Linn Co., IA

Bailey

Nutting 1892, Anderson 1907

Record Number: 81-HJ  
Classification: NA

REFERENCES

Nutting, C.C. 1892. Report of the committee on state fauna.  
Proc. Iowa Acad. Sci. 1(Part 3):39-42.

VOTE (1981): 4-IV, 3-V

IV, no data.

IV, unclear who saw this bird.

IV, no detail; again, probably correct at that time.

Nutting, C.C. 1892. Report of the committee on state fauna.  
Proc Iowa Acad Sci 1(Part 3):39-42.

*Calcarius ornatus*, (Townsend).—~~Chestnut-collared Longspur~~ Cedar Rapids, Iowa (Bailey.) ] 6

Anderson 1907  
p 314

237. (538). *Calcarius ornatus* (Townsend). Chestnut-collared Longspur.

The Chestnut-collared Longspur is a bird of the western plains, occurring in Iowa rather early as a migrant or winter visitant. The species is not known to breed in Iowa, although in 1896 J. H. Brown and J. Eugene Law found two nests in Jackson county, Minnesota, a short distance north of the Iowa line.

County records: Howard—"more common than any other species in winter. They are, I believe, the most common bird breeding in South Dakota. Do not breed here" (E. B. Webster, Cresco). Jefferson—"I identified this bird in Jefferson county, Iowa, in 1896" (W. G. Savage). Linn—"Cedar Rapids, Iowa, —Bailey" (Nutting, Proc. Iowa Acad. Sci., 1892, p. 40); "common winter visitant" (Berry). Poweshiek—"rare winter visitant" (Kelsey). Pottawattamie—"common migrant" (Troster). Van Buren—"winter resident; very rare" (W. G. Savage). Winnebago—"shot at Forest City in 1903" (Halvorsen).

4  
2  
] 5  
7  
8  
9  
3

DuMont 1933  
p 154

*Calcarius ornatus* (Townsend). Chestnut-collared Longspur

A fairly rare migrant in the western part of the state. Anderson (1907) listed it from Howard, Linn, Van Buren, Poweshiek, Jefferson, Pottawattamie, and Winnebago counties. He stated that M. Earle Halverson shot one at Forest City, Winnebago County, in 1903.

Lynds Jones recorded (Wilson Bull., IV, p. 27) that during 1887 several hundred remained for several weeks at Grinnell, Poweshiek County. Spurrell (1919) recorded two seen at Wall Lake, Sac County, during the fall of 1912. Fenton (1923-24) recorded that a specimen taken at Charles City, Floyd County, and contained in the Miles collection, was identified as this species by B. H. Bailey. Wolden reported that he observed a large flock of these birds in Emmet County late in the winter of 1932-33. He wrote that he was able to make out the whitish throats and the dark breasts, although the birds were not in full breeding plumage. The writer has examined no Iowa specimens.

✓  
1  
10  
11  
12