Records Committee, Iowa Ornithologists' Union Printed: 09/07/93

Painted Bunting 6 Jan 1987 Bettendorf, Scott Co., IA *P. Petersen, *A. Barker IBL 57:57, 58:76, P-0218 Record Number: 87-06 Classification: A-E

DOCUMENTATIONS

Peter C. Petersen Ann Barker PHOTOGRAPHS P. Petersen, P-0218 REFERENCES

Field Reports: IBL 57:57 Records Committee: IBL 58:76

VOTE: 2-I-P, 2-III, 1-V, 1-VI, 1-no vote

III, I hope the photographs are conclusive, because notes were not written for five hours or four days after the observation. For a potential first state record, this is not very encouraging. The only question is whether this bird was a naturally occurring vagrant. There is a male of this speices at a feeder in Nebraska at just about this same time. I heard about it on a rare bird alert, so don't know the details or outcome of that record.

V, Adult males have same plumage year around, although it may not be attained until second winter. Midwest (NE, MN, WI, IL, IN, MI, ONT) records are distributed as follows: 5 in Apr, 12 in May, 2 in Jun, 1 in Sep, 1 in Oct, 2 in Nov. East Coast records (NB, ME, NH, MA, RI, NY, NJ, PA, MD, VA) are distributed as follows: 4 in Jan, 5 in Apr, 28 in May, 3 in Aug, 2 in Sep, 1 in Oct, 2 in Nov, 4 in Dec. Bent describes how males were easily caught and common as cage birds. Roberson says it has been illegal to hold these birds since the early seventies. Almost all Nov-March birds have been present several days to several weeks. The date of this sighting and fact that the bird did not stay around are very unusual. The plumage does not argue one way or another. The first Iowa record, although undescribed, is a better record because the date and location argue against an escapee and the competent observers could hardly misidentify it.

no vote, I have no doubt what was seen and described was a Painted Bunting. The descriptions are not even close to anything else. I tend to look at this record as a possible escapee rather than a highly likely winter vagrant. This species is colorful enough and adaptable to caging that this possibility cannot be ruled out, especially in mid-winter. This species should be 1000+miles south of Iowa at this time of year. Since Scott Co. is not a great migrant trap like the East Coast tends to be I am extremely cautious about the origin of this particular bird.

VI, I have no doubt that the bird was a painted bunting but I need more input on vagrancy patterns of this species at this unlikely time of year. The bird may have been an escaped bird to be here in Jan.

I-P, I have seen photo. While possibility of escape exists, there is precedent for vagrancy in this species. I believe this is a record of a wild bird. However neither observer addressed

Records Committee, Iowa Ornithologists' Union Printed: 09/07/93
Painted Bunting 6 Jan 1987 RC No. 87-06 (cont)

this issue.

III (I-P with photos), Bizarrre. Only real question is whether it might perhaps be an escape.
REVOTE (at meeting, 30 May 1987, with photos): 4-I-P, 2-V, 1-abstain

For Extraordinary Bird Sightings in Iowa

87-06 PG

What species? Tainted Bunting of How many?
Location? 1213 Broadlayn, Bettendorf, Scottle # A.
Type of Habitat? urban yard near woods
When? date(s): 6 Jan, 1987 time: 8:00 AM to to 9:15P. Moff Who? your name and address: Peter C. Petersen, 235 McClellan Blud, Davenport, IA5.390
Who? your name and address: Peter C. Petersen, 235 McClellan Blud, Davenport IASson
Others with you: Bety Hunt, Ann Barker Gil Sandvick, Brian+Corey Blevins
Others before or after you: <u>before-leah Leviu-</u>
Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, undertail, legs, feet). Also mention voice and behavior. Fairly small finch, smaller than nearby thouse 5 perrous x slightly larger than 5 hins, same general 5 he pe as House 5 perrous, throat, breast, belly + under fail as well as rump+lover back bright red-orange more to red than orange. Remainder of head bright blue like top of more to red than orange. Remainder of head bright blue like top of head of Fadigo Benting except evering red-orange, Back & upper uing head of Fadigo Benting except evering red-orange, Back & upper uing coverts reliangemen. Using satail brown, Bill conecal, grey, lighteron (over mandible; legs darkgrey, Noth eard, worked below niger feeder and 5 at in & below nearby bushes, Associated with Mcardinals. See photos.
Similar species; how eliminated: Nothing is similar in any way,
Did any one disagree or have reservations about the identification?w ©
Viewing conditions: give lighting, distance (how measured), optical equipment. Bird within 15-18' Tun Fromright (X40 binoculars, 5x+20x Lenses
Previous experience with species and similar ones: Have seen frequently in Fexas + Ollahouna.
References and persons consulted before writing description: none needed.
How long before field notes made? 5hv5 this form completed? 5hv9.
Send completed form to Field Reports or CBC editor (address on back cover of Iowa Bird Life).

DOCUMENTATION FORM
For Extraordinary Bird Sightings in Iowa

What species? <u>Painted Bunting - male</u> How many? 1
Location? 12/3 Broadlawn, Bettendorf Ja.
Type of Habitat? Under and in vicinity of backyard feeder - in nearly bushes
When? date(s):
When? date(s): Jan 6, 1989 time: 9 AM to 9:15 Who? your name and address: Ann Barker, Rl3, Box 190, Dovenport, La
Others with you: Pete Petersen, Brian & Corey Blevius, Leah Levin, Betty Hunt, Gil Sanderck
Others before or after you: _?
Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, undertail, legs, feet). Also mention voice and behavior.
Gaudy Donabird - angray the sing of a Long Sparrow or Purple Time
Gaudy Dongbird - approx the size of a Long Sparrow or Purple Finch. Finch like bill. Head blue, with red eye ring. Breast, belly & rump
all bright rid. Back chartreuse-green. Wings and tail brownish
with hint of green. Fed on ground beneath feeder; flew to rest
with hint of green. Fed on ground beneath feeder; flew to rest in bushes or on ground 15-20 feet from feeder.
Similar species; how eliminated:
Did any one disagree or have reservations about the identification?
Viewing conditions: give lighting, distance (how measured), optical equipment. Cloudy, but decent light. Seen through window at destances of 10-30 feet. Lette 10x40 binoculars
Previous experience with species and similar ones: None
. References and persons consulted before writing description: Nat. Goog. Givede, Peterson + Robbins field quedes (4 days)
How long before field notes made? 1-10-87 this form completed? 1-10-87
Send completed form to Field Reports or CBC editor (address on back cover of Iowa Bird Life).