

Season: Fall Year: 82Name and address
(print or type) M Newton

Other observers (name, city):

Use first few lines to give weather and habitat conditions. Use line(s) below species data to give additional details, if needed. Mail to:
T. H. Kent, 211 Richards St., Iowa City IA 52240
by 1 Mar, 1 Jun, 1 Aug, or 1 Dec.

use initials of obsv. on species line

SPECIES	NO.	DATE	LOCATION - COMMENT - OTHER OBSERVERS
			(all records Coalville Res. area)
✓ pelican W	15	10-17	Co Res
d-cormorant	150	10-17	" " max; present 9-19- 10-31
gt blue heron	90	9-19 to 10-23	
gt egret	20	9-19	" max
cattle egret	15	9-19	Tom's Pond Johnson Co
rb grebe	500	11-27	max; present 10-17 on
✓ t vulture	30	8-1	
✓ osprey	1	9-27	
bald eagle	1i	10-22	
	3a	11-27	
✓ n-1 hawk	1	10-22	1st record
✓	1	11-27	(Iowa City)
✓ peregrine	1	9-27	
✓	1	10-10	(different bird)
✓ merlin	1	10-10	(photo)
✓ o-s flycatcher	1	9-19	
e phoebe	1	10-3	
✓ water pipit	2	9-27	
✓	2	10-17	

Shnebuds - S. side Co. Res. 1982

	7.31	8.21	8.28	9.12	9.19	9.27	10.3	10.10	10.17	10.23	10.31	11.7
serripal. pl.		20	20	10	30	12	5					
kilclen.	10		10	6	2		13	3	9	14	17	2
l. golden pl.		6		6			2					
bl-b. pl.						3						
mudch + s				1								
gr. y-l.					8			10	14	5	4	15
le. y-l.	30	40	50	50	30	35	27	5		3		1
sol. s.	10		1									
spot. s.	6											
W. phal.	1i				1i							
N. "			1									
C. Snipe					5		4	1	48	35	32	8
sb dow		3i	1i									
l-b dow						2i		1i	2i		1b	1b
^{sp} sanderling					1	5		7		3		
semi. s.		3	30	25	30	45	6	1	11	15		
W. s.						2	2					
lot. s.	3	30	1	3	4	70	5		3			
Band's s.						1i	2(ii)			1		
pect. s.	3	50	200	300	600	60	355	27	89	93	66	4
dunlin						1	1		6	30		
stilt. s.			22i	30i	20i	6			1	1		

descriptive details

S-b juv - in full plumage, with tiger-striped tertiaries very obvious.

I-b - same birds present throughout - most were at least partially molted into Basic plumage; but ~~not~~ on all birds identified as long-bills at least a few of the characteristic juv. scapulars on tertiaries were visible, dark with narrow rusty edges and no internal markings.

dr. teta details

S-b juv - in full plumage, with tiger-striped
tertials very obvious.

l-b - same birds present throughout - most were
at least partially molted into basic plumage;
but ~~not~~ on all birds identified as long-bills
at least a few of the characteristic juv.
scapulars on tertials were visible, dark with
maroon rusty edges and no internal marking.