Records Committee, Iowa Ornithologists' Union Printed: 09/01/93

Black-legged Kittiwake
21 Nov 1984
Saylorville Res., Polk Co., IA
*D. Thompson, *M./*B. Proescholdt, *Moore
IBL 54:115; 55:57

Record Number: 84-29 Classification: A-D

DOCUMENTATIONS

Dennis Thompson (21 November) Beth Proescholdt (21 November) Mark Proescholdt (21 November) Francis L. Moore (22 November)

REFERENCES

Field Reports: IBL 54:115 Records Committee: IBL 55:57

VOTE: 7-II

II, Excellent descriptions.

II, Well described. Good sketch by Moore.

II, Note: the first documentation didn't copy legibly. Probably doesn't matter in this case. Francis Moore's documentation is excellent.

(Iowa, Missouri, Illinois, Indiana, Kentucky, Ohio) BIRDS: VERIFYING DOCUMENTATION OF AN EXTRAORDINARY SIGHT RECORD. 1. Species Black-leaged Kittiniake 3. Location Selow dam at Saylorville Pes. Polt Co. Fowa 5. Time Bird seen: 10:00 : 6. Description of size, shape and color-pattern (describe in great detail all parts of the plumage, and beak and feet coloration, in addition, to the diagnostic characteristics, but include only what actually was seen in the field): Kird was slightly smaller than the Ring-billed gulls itwas feeding with. It was pale-bodied with a black half-collar. There was a block spot half-way between the collar and the eyes. There was a dark "w" across the wings (and back!) The primaries were all-dark. The back and is of the upper wing was gray. The bill was small, black thip! The legs were bluck. The tail was slightly forked with an obvious black band. 7. Description of voice, if heard: 8. Description of behavior: Resembled a Bonaparte's Inflight. Would plunge into the water 9. Mabitat - general: outlet below the dam. Feeding in the fast-moving water directly was the dam. 10. Similarly appearing species which are eliminated by questions 6, 7 & 8, Explain: Similar in shape actions to Bunaparte's, No direct comparison was available, but somed larger I'd ray half-way in size between Bonaparte's + Ring-billed. Black legs, black half-collar, and lack of carpal bar helped eliminate Bongpartes, Coray back took care of subino's, white heady Franklin's. 11. Distance (how measured)? Cloce range - less than 304ds, 12. Optical equipment: Bros, 13. Light (sky, light on bird, position of sun in relation to bird and you): Bright sun behind me, 14. Previous experience with this species and similarly appearing species: Observed one at Superville in 1981. Familiar with Bongparte's, 15. Other observers: by my celf, 16. Did the others agree with your identification? Other observers who independently identified this bird: Edgene + Eliste Armstrong, Beth+ Mark Preschielt, Dick+ Pauly Mooney, Marycret Brooks 18. Books, illustrations and advice consulted, and how did these influence this description: Had Robbins Browk with me, Looked at Harrison's, Nat. Ceu, hooks later. 19. How long after observing this bird did you first write this description? / week. Address: 2019

City, State:

84-29 DOCUMENTATION FORM for extraordinary bird sightings in Iowa What species? Black-legged Kittiwake, imm. How many? / Location? Saylorville Reservoir Type of habitat? below spillway of dam When? date(s): November 21, 1984 time: 12145 to 12150 P.M. Who?your name and address: Beth Proescholdt, Liscomb, Iowa others with you: Mark Proeschold + Eloise + Gene Armstrong, faully + Diskney, others before or after you: Denny Thompson, Francis Moore Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, under tail, legs, feet). Also mention voice and behavior. A medium-sized gull, smaller than the Ring-bills flying around it, this bird with soft gray mantle + white body had inky black wingtips with the black continuing on from the wrist slanting across the secondaries to form an M pattern across the upperwings. It had a dark bar across the nape, a dark eye with a black spot behind the eye, a black bill and a narrow black band across the end of the tail which was slightly torked, but seemed more so by the indented black band; warming The few minutes we saw it the bird was flying and feeding above the water just below the spillway, we observed it about 5 minutes after which it flew up + over the dam, + we didn't see it again in our hour of waiting. Similar species and how eliminated: immature Bonaparte's Gull; - this Kittiwake had no white in its wingtips as a Bonaparte's, + it had a black bar on its nape. The tail appeared forked while a Bonapartes is slightly rounded. Did any one disagree or have reservations about identification? No If yes, explain: Viewing conditions: give lighting, distance (how measured), and optical equipment: binoculars - from parking lot we looked down on gull

MAIL TO: T. H. Kent, Field Reports Editor, 211 Richards Street, Iowa City IA 52240

Previous experience with species and similar ones: Never a Kittiwake, but several

immature Bona parte's
References and persons consulted before writing description: Guides; Peterson's, Robbins,
National Geographic, Audubon Soc. Master Guide.
How long before field notes made? <u>At once</u> this form completed? <u>Nov.</u> 23,1984

	DOCUMENTATION FORM for extraordinary bird sightings in Iowa 84-29
	what species? Black-legged Kittiwake How many? 1
	Location? Just below Saylorville Dam - (Polk Co.)
	Type of habitat? Churning water below reservoir dam in large river valley when? date(s): November 21, 1984 time: 12:45 to 12:50 p.m very valley
	When? date(s): November 21, 1984 time: 12:45 to 12:50 p.m very approximate time
	Who?your name and address: Mark Proeschold T Liscomb, Iowa
	others with you: Beth Proescholdt, Gene and Eloise Armstrong Margaret Brooks others before or after you: Dennis Thompson - before, Francis Moore-after
	others before or after you: Dennis Thompson - before, Francis Moore-after
	Describe the bird(s) including only what you observed. Include size, shape, details of all parts (bill, eye, head, neck, back, wing, tail, throat, breast, belly, under tail, legs, feet). Also mention voice and behavior.
	An immature gull was swooping low over the churning
	voter below Saylorville Dam. It had a white head with a
1	plack cheek spot and a dark bar across the back of its neck.
	It had black wing tips and very dark bars across the inner wings.
,	the tail was slightly forked with a black tail band. I did not see
,	the legs on this bird.
	It was flying by itself, but it seemed bigger than the Bonaparte's gulls we had seem and work and in Tigger than the
	Bonaparte's gulls we had seen one week earlier, It was swooping
	low over the water. Then it gained elevation, swirled around
	a bit, and sailed over the dam. It was not seen again by me
	Similar species and how eliminated: An immature Bonaparte's Full does not have the black bar on the nape and has white in the outer primaries.
	Did any one disagree or have reservations about identification? No
	If yes, explain:
	Viewing conditions: give lighting, distance (how measured), and optical equipment: It was a sunny, clear day with the sun behind us. Distances
	estimated. 7x35 Binoculars.
	Previous experience with species and similar ones: First sighting of a Kittiwake. Saw quite a few immature Bonaparte's gulls at Saylorville one week before. References and persons consulted before writing description: Natl. Geog. Birds of N. America, R.T. Reterson's Birds East of the Rockies, Robbins' Birds of N. America.
	How long before field notes made? Right after this form completed? Nov. 23, 1984 viewing bird. MAIL TO: T. H. Kent, Field Reports Editor, 211 Richards Street, Iowa City IA 52240
	TIME TO. 1. II. Kelle, Fleta Reports Editor, Ett Kichards Solices, Iona Grof In Section