Records Committee, Iowa Ornithologists' Union Printed: 08/19/93

Black-backed Woodpecker 23 Dec 1978 north of Boyden, Sioux Co., IA *Jay A. Stravers IBL 49:26,56 Record Number: 81-CN Classification: A-D

DOCUMENTATION

Jay A. Stravers

REFERENCES

Field Reports: IBL 49:26, 56

VOTE (1981): 7-III

III, bare minimum detail, behavior fits.

III, exemplary documentation.

III, OK description in documentation from 3 observers.

III, 'Meticulous' documentation accepted by Halmi and appears sound to me.

Erchiesestern grairie Regisa (Iowa, Missouri, Illinois, Indiana, Pentucky, Ohio) = Grie 81-CN RECORD. Since BIRDS: VERIFYING DOCUMENTATION OF AN EXTRAORDINAL Speck Backed Three Toed Wood pecker, Picoides arctize. Number: Location 21/2 miles North of Boyden Jours on east side of county highway 14 5. Time Bird seen: 1:00 PM to about 2:30 PM temp. - 4°F winds from the N-NW 10-20 mph relatively low barametric pressure (less than 30.0am) Description of size, shape and color-pattern (describe in great detail all parts of the plumage, and beak and feet coloration, in addition, to the diagnostic characteristics, but include only what actually was seen in the field): ic - body size slightly larger than heiry woodpecker, around 25cm from crest to tip of tail (quesstimate) d + body shape rather like hairy woodpecker. Beak - black - 4-5cm long. Feet - very dark almost black to stock each about 3-4cm long. Logs invisible. Fewfureless black back. Wings black with white specks. er tips of wings, no discernable pattern to these white specks because plumage was ruffled by winds. uture less white breast. Short (less then Icm.) black bars on the bird's sides. Bars are roughly rpen dicular to the long axis of the birds body. Throat is white with a thin black stripe on both. The running dem from the base of the beak down both sides of the throat. There is a bright yellow the covering a nearly the entire crest but not appearing on the back or sides of the birds headso it invisible if you were looking up at the bird at an angle greater than about 45° measured upmand from horizon. Eyes black with no eye rings. Short white whishers at the base of the beak. OVER Description of voice, if heard: no voice heard possibly feeding? Description of voice, if heard: NO voice heard

This print berk from trees (mostly dead elms). The birds spent an average of the Description of behavior: about 5 mins. at each individual tree, usually starting near the base of the tree and working upwards to a height that never exceeded about 10 m. Size of bark chips varied up to about 25cm² behavior of bird could have been effected by Proximity of observers, but the bird seemed totally unwary antill approached to within 3 m. Habitat - general: 3000 m farm grove in wide open (formerly prairie) gently rolling farm land. specific: grove is a thick growth of shrubs, weedy plants, Ash trees, and American + Chinese Elms. Tany of the Elms are cleud and fallen. There are 2 rows (closely spaced) of Ponderosa Pine on the North side of the grove. The grove is sucrounded with Mature Honory such the Shrubs. Grood covered with 0.5 m. of such such appearing species which are eliminated by questions 6,7 & 8, Explain: in the grove. PARTO BESTORE There are really no other species that could be confused other than possibly a Pileated Woodpecker who is much too large + has 4 toes instead of 3 Distance (how measured)? 12. Optical equipment: naked eye, 8 x 50 Binocular, 60mm Bushnell specementer - 20 x exercice 1.50 meters to 30 meters (estimated) Light (sky, light on bird, position of sun in relation to bird and you): recast sky. Bird was observed from virtually every different sun angle Previous experience with this species and similarly appearing species: no previous experience with this species. I am well acquainted with the Downy, Hairy, and Other observers: Fileated Loreen Stravers, Honry O. Stegeman, Aaron Stegeman Pileated woodpukers Did the others agree with your identification? yes Other observers who independently identified this bird: Lorcen Stravers Books, illustrations and advice consulted, and how did these influence this description: Robbins, Brunn, Zin, and Singer; 1966; Birds of North America Peterson - A Field & Guide to the Birds. The Audubon Society Field Guide to North American Birds (Western Region) scription of the bird given in 6. was rewritten from field notes taken during observation period books were used only for identification + general information. They did not influence the lover Clay a. Stravers Jay A. Stravers Address: 3220 Madison Ave #2 nature. Boulder City, State: Colorado 80303 e. 5-13-29 estions - this form lacks on entry for specific weather conditions at time of sighting

nt.

I mostly black about 8cm long. There are some white feathers in the tail but there iten in the tail could not be determined because of wind ruffled plumage.

Halmi, N. S. 1979. Field reports. Iowa Bird Life 49:23-26 (p 26)

scarce. On December 23, Mr. and Mrs. Jay Stravers saw a male Black-backed Three-toed Woodpecker near Boyden, Sioux Co. (fide GB). Since there is no recent record of this species in the state, documentation is eagerly awaited.

Halmi, N. S. 1979. Field reports. Iowa Bird Life 49:56-63 (p 56)

Jay Stravers furnished meticulous documentation for the male Black-backed Three-toed Woodpecker he, Mrs. Stravers and two friends studied at leisure and from as close up as 1.5 m on December 23, 1978. This is the first Iowa record since 1924, and only the third time the species has been seen in Iowa. A Varied Thrush at a feeder in

tion other than to suggest particular details to observe note while making observations