Lougerbead Shrike--The pair at LaTROPREPORTEL Destrong five young on June Multure

Bell's Vireo--Three located, two at Sweet Marsh June 17, one at Volga Lake July

Pied-billed Grebe--Few found in the northeast, with the exception being 15 at Cardinal Marsh, Winneshiek Co., on June 18, most being immatures. Green Heron--Down in numbers, suggested to be watched.

Least Bittern--Two or more were found in June at Cardinal Marsh, during each visit, Green-winged Teal--Two males were at Sweet Marsh, Bremer Co., on June 14. Lesser Scaup--One female was at Cardinal Marsh on June 12. Swainson's Hawk--Nest at Fayette Co. appeared successful, with one young last seen there attempting to fledge July 15. An Adult bird was seen near Alpha, Fayette Co., on July 7, and may have bred in area.

Common Bobwhite--Abundant in Volga Lake, but absent from the rest of Fayette Co., showing what a little suitable habitat can do.

Wild Turkey--A nest was found in Volga Lake, Fayette Co., by Luther College people during late june. The population at Volga Lake is good.

Virginia Rail--An obtrusive bird at Sweet Marsh on June I4, was probpably protecting young.

Semipalmated Plover -- A late bird was at Sweet Marsh on June I4.

Upland Sandpiper--A total of I3, were seen in northeast Iowa (Chickasaw, Fayette, Howard, and Winneshiek) counties, June 9 to July I3. The highest single flock was

- that of four near Fredricksburg in Chickasaw Co., on July 9. Woodcock--This uncommon bird was rather common in Volga Lake this summer. Common Snipe--Appearance of this species twice in June is suspicious. One was found
- at St. Lucas on June I8. One was at Sweet Marsh June I9. Black Tern--A flock of I5 was at Cardinal Marsh on June I2. Ruby-throated Hummingbird--Fairly common at Volga Lake during June. Belted Kingfisher--Scarce along northeast rivers, no improvement seen. Eastern Phoebe--Rather common in Volg Lake, uncommon in rest of county. Acadian Flycatcher--Four plus again summered at St. Lucas, and one or two were found in Volga Lake on June 29 for a new location.

Alder Flycatcher--Four were located during period in northeast; one five miles northeast of New Hampton on June 13; three in Volga Lake on July 6, see general notes write-up for details.

Least Flycatcher--One was regularly found into July at Volga Lake.

Tufted Titmouse--Only three could be located during period in Fayette Co. Brown Creeper--This species was located at Sweet Marshoon June I7, which was an adult feeding a recently fledged young.

Veery--Two were found regularly at Sweet Marsh into late June. Eastern Bluebird--Still slipping and may be extripated from most of Iowa.

-OVER-

Loggerhead Shrike--The pair at Lawler had a nest containing five young on June 24, a pair was also located south of Alpha.

Bell's Vireo--Three located, two at Sweet Marsh June I7, one at Volga Lake July 6. Blue-winged Warbler--Thirteen were located in late June in Volga Lake so is doing well there.

Chestnut-sided Warbler--One was at Sweet Marsh on June I4, and six were found in Volga Lake in late June and early July, see general notes write-up. Eastern Meadowlark--Abundant in Volga Lake, uncommon or in small numbers elsewhere. Orchard Oriole--One or two was found on June30 in Volga Lake.

Swainson's Hawk--Nest at Fayette Co. appeared successful, with one young last seen there attempting to Clader Wild; purch puilaged near Alpha, Fayette Co., on July 1914 and have breding reaching in puilage

Common Bobwhite Moundant in Volga Lake, but absent from the rest of Fayette Co., showing what a little suitable habitet can to. Wild Turkyv-4 Mat was file a sign of the file, Fayette Co., by Luther College people

during an der in population at Volge Le is good.

Virginia Rail--An obtrusive bird at Sweet Marsh on June 14, was probably protecting young.

Semipalmated Plover--A late bird was at Sweet Marsh on June 14. Upland Sandpingr--A total of 13, were seen in northeast Iowa (Chickasaw, Fayette, Howard, and Winneshiek) counties, June 9 to July 13. The highest single flock was that of four near Fredricksburg in Chickasaw Co., on July 9.

Woodcock--This uncommon bird was rather common in Volga Lake this summer.

Common Snipe--Appearance of this species twice in June is suspicious. One was found at St. Jucas on June IS. One was at Sweet Marah June I9.

Black Tern--A flock of I5 was at Cardinal Marsh on June I2. Ruby-throated Hummingbird--Fairly common at Volga Lake during June. Belted Kingfisher--Scarce along northeast rivers, no improvement seen. Eastern Phoebe--Rather common in Volg Lake, uncommon in rest of county. Acadian Flycatcher--Four plus again summered at St. Lucas, and one or two were found in Volga Lake on June 29 for a new location.

Alder Flycatcher--Four were located during period in northeast; one five miles northeast of New Hampton on June 13; three in Volgs Lake on July 5, see general notes write-up for details.

Least Flycatcher -- One was regularly found into July at Volga Lake.

Tufted Titnouse --- Only three could be located during period in Fayette Co. Brown Greeper--- This species was located at Sweet Marshoon June 17, which was an adult feeding a recently fledged young.

Veery--Two were found regularly at Sweet Marsh into late June. Restern Bluebird--Still sliping and may be extrapated from most of love.